

An aberrant ‘‘spin-orbit interaction’’ persists in the literature since more than thirty years.

Jacques Raynal

4 rue du Bief, 91380 Chilly-Mazarin, France

Abstract

An expression for the spin-orbit interaction coupling between different levels, which was shown to be aberrant more than thirty years ago is used in a recent article [1] published in Nuclear Physics. It leads to expressions quite simpler than they should be. Its behavior is in fact of a character opposite to that of the spin-orbit interaction used in two-body studies.

PACS numbers 24.10.-i, 25.40.Dn, 25.40.Ny, 28.20.Cz

In the article entitled “An algebraic solution of the multichannel problem applied to low energy nucleon-nucleon scattering”, written by K. Amos, L. Canton, G. Piseni, J. P. Svenne and D. van der Knijff, published in Nucl. Phys. **A728** (2003) 65–95, the authors describe in their Appendix B the interaction which they use. On the second line of page 92 :

$$-\frac{1}{2\alpha r}W_{ls}\{[\ell.\mathbf{s}]_{c'} + [\ell.\mathbf{s}]_c\} \quad (1)$$

with the two following lines deals with the spin-orbit interaction. This is a generalisation of the spin-orbit potential of the optical model, which is :

$$\frac{2}{r}\left\{\frac{d}{dr}V(r)\right\}[\ell.\mathbf{s}] \quad (2)$$

and was used when first asymmetry measurements in inelastic scattering with polarised proton beams became available. The expression Eq. (1) was used in [2] without being explicitly written in this article. It was sometimes called Oak-Ridge term and gave results quite different from the experimental results. Some attempts [3] were done to get better results, adding ℓ^2 and $\mathbf{s}.\mathbf{I}$ terms which are also included in [1].

Going back to the “full Thomas term” obtained for the spin-orbit by transforming a Dirac equation into a Schrödinger equation, J. S. Blair and H. Sherif [4, 5], used the expression :

$$\nabla\{V(\mathbf{r})\} \times \frac{\nabla}{i} \cdot \boldsymbol{\sigma} \quad (3)$$

in computations for nucleon inelastic scattering and obtained a very good reproduction of the experimental data.

This expression can be written [6, 7] like Eq. (1). Using $V_\lambda(r)$ for the radial dependence of a multipole λ , the result is :

$$\begin{aligned} \frac{1}{r} \frac{d}{dr} V_\lambda(r) [\ell \cdot \mathbf{s}]_c + \frac{V_\lambda(r)}{2r^2} \left(\lambda(\lambda+1) - ([\ell \cdot \mathbf{s}]_{c'} - [\ell \cdot \mathbf{s}]_c)([\ell \cdot \mathbf{s}]_{c'} - [\ell \cdot \mathbf{s}]_c \pm 1) \right) \\ + \frac{V_\lambda(r)}{r} ([\ell \cdot \mathbf{s}]_c - [\ell \cdot \mathbf{s}]_{c'}) \frac{d}{dr} \end{aligned} \quad (4)$$

with $+1$ or -1 if the wave function is not or is multiplied by r . Note that there are here three form-factors :

- $\frac{1}{r} \frac{d}{dr} V_\lambda(r)$ which is the only one for elastic scattering and is multiplied only by the eigenvalue for the ket.
- $\frac{V_\lambda(r)}{2r^2}$ which is, divided by r^2 , the true spin-orbit multipole which does not appear in elastic scattering.
- $\frac{V_\lambda(r)}{r} \frac{d}{dr}$ which is the form factor multiplying the derivative of the ket radial function; integrating by part shows that the whole is symmetric in c and c' .

Except for the first, the form factors are not the ones of Eq. (1). However, as the interaction of Eq. (1) is larger when the eigenvalues for c and c' are of the same sign and the coefficients of the second and third form-factors above are larger in the opposite case, one can guess that the effect should be quite different. The use of the spin-orbit deformation given by Eq. (4) in coupled channel calculation is not straightforward [8]. It was the subject of codes ECIS (“*Equations Couplées en Itérations Séquentielles*”) from ECIS68 to ECIS97. To compare results obtained with the interactions given by Eq. (1) and Eq. (3), the spin-orbit interaction is parametrised as :

$$\begin{aligned} \frac{1}{r} \frac{d}{dr} V_\lambda(r) \left(z_1 + z_3 [\ell \cdot \mathbf{s}]_c + z_4 [\ell \cdot \mathbf{s}]_{c'} \right) + \frac{V_\lambda(r)}{r} z_6 ([\ell \cdot \mathbf{s}]_c - [\ell \cdot \mathbf{s}]_{c'}) \frac{d}{dr} \\ + \frac{V_\lambda(r)}{2r^2} z_5 \left(z_2 \lambda(\lambda+1) - ([\ell \cdot \mathbf{s}]_{c'} - [\ell \cdot \mathbf{s}]_c)([\ell \cdot \mathbf{s}]_{c'} - [\ell \cdot \mathbf{s}]_c \pm 1) \right) \end{aligned} \quad (5)$$

in all these codes. The coupling of Eq. (1) is obtained by setting :

$$z_1 = z_2 = z_5 = z_6 = 0, \quad z_3 = z_4 = \frac{1}{2}, \quad (6)$$

the coupling of Eq. (3) multiplied by a parameter λ (with $\lambda = 1$ for the “unparametrised” case) is given by :

$$z_1 = z_4 = 0, \quad z_2 = 1, \quad z_3 = z_5 = z_6 = \lambda. \quad (7)$$

The parameter λ allows to increase the strength of the spin-orbit transition without deforming its form factor in the rotational model. The use of $\lambda = 2$ for experiments around 20 MeV gave excellent results.

For people who don’t want to consider Dirac equation at low energy, there is another justification of Eq. (3) based on the nucleon–nucleon interaction [6, 7, 9]. At the zero-range limit, the two-body spin-orbit interaction of a nucleon exciting a particle–hole state is given by Eq. (4) with the product of the particle and the hole functions as V_λ , assuming that the sum of the eigenvalues of $\ell \cdot \mathbf{s}$ for the particles and holes vanishes in the result. The overestimation by a factor 4 of the spin-orbit interaction in the first publications on this subject is an error which does not affect this similarity. This approach, with the most general consideration of the two-body interaction [10] has been the subject of a series of codes, since DWBA70 to DWBA98. Note that spin-orbit interaction involves first derivatives, quadratic spin-orbit involves second derivatives. In this approach, the geometrical coefficients are expressed with $\kappa = [\ell \cdot \mathbf{s}] + 1$ which is the coefficient of $\frac{1}{r}$ in Dirac equation. For a central interaction $V(r)(\boldsymbol{\sigma}_1 \cdot \boldsymbol{\sigma}_2)$ and a tensor interaction, the geometrical factor can be reduced to the one involved by a spin-independent interaction leading to two one-body form-factors : one simple and one multiplied by the sum or the difference of the κ . Usual and quadratic spin-orbit need more form-factors. The two-body interaction which keeps the parity for each body is characterised by coefficient $\kappa_c - \kappa_{c'}$; the one which inverts them is characterised by $\kappa_c + \kappa_{c'}$, equal to a constant with what appears in Eq. (1) but has no equivalent in one-body interaction (the spin-orbit has no derivative but a third form-factor with $(\kappa_c + \kappa_{c'})_2$).

The reference [10], in which the first co-author is the same as for [1] shows that some users of DWBA or ECIS do not realise what these codes involve. In fact, the author of this “comment” cannot answer on many points of the article [10].

There is no allusion to a second matrix with derivative of the ket $|c\rangle$ in [1]. In fact, its Eqs (36) and (42) includes expressions like $[\ell \cdot \mathbf{s}]_{c'c}$ which could be a shorthand notation of a correct spin-orbit interaction; this cannot convince the reader that the formulae of Appendix B are not used. The model presented here is the two-phonons vibrational model limited to one kind of phonons; there is no factor $\kappa_c + \kappa_{c'}$ which appears only for two phonons of angular momentum L_1 and L_2 coupled to a J such that $L_1 + L_2 + J$ is odd.

The Eq. (1) is often written by authors to say that they do not use it and cannot be found in [2, 3]. We cannot know how many works used it since thirty years; the main interest of [1] is to show that it continues to be used by some physicists, at least from time to time. Its behaviour being opposite to the one used in two-body studies and its lack of justification (except the great simplification of the problem) justify the qualification of “aberrant”.

I thank the Service de Physique Théorique de Saclay for allowing me to follow this kind of problems after my retirement. I also thank H. Sherif for

reading the manuscript and for helpful correspondence.

References

- [1] K. Amos, L. Canton, G. Piseni, J. P. Svenne and D. van der Knijff, Nucl. Phys. **A728**, 65 (2003).
- [2] T. Tamura, Rev. Mod. Phys. **37**, 679 (1965).
- [3] O. Mikoshiba, T. Terasawa, and Tanifugi, Nucl. Phys. **A168**, 417 (1971).
- [4] H. Sherif, J. S. Blair,, Phys. Let. **26B**, 489 (1968).
- [5] H. Sherif, Spin-dependent effects in proton inelastic scattering, Thesis University of Washington (1968).
- [6] J. Raynal, Symposium sur les Mécanismes des Réaction Nucléaire et les Phénomènes de Polarisation, Québec, Canada, Sept.1-2, 1969, Les Presses de l'Université Laval, Québec, 75 (1970).
- [7] J. Raynal, The Structure of Nuclei, International Course on Nuclear Theory, Trieste, Italie, Jan.13 - March 12, 1971, (IAEA,1972) 75.
- [8] J., Raynal, Computing as a Language of Physics, ICTP International Seminar Course, Trieste, Italy, Aug.2-10, 1971
- [9] J. Raynal, Nucl. Phys. **A97**, 572 (1967)
- [10] K. A. Amos, P. J. Dortmans, H. V. von Geramb, S. Karataglidis, J. Raynal, Adv. Nucl. Phys. **25**, 275 (2000) (IAEA, 1972), 281.